

## Harland MCB 231 2007 Xenopus

### Reviews

[Harland, R.](#) (2000). Neural induction. *Curr Opin Genet Dev* 10, 357-362.

\*[Harland, R., and Gerhart, J.](#) (1997). Formation and function of Spemann's organizer. *Annu Rev Cell Dev Biol* 13, 611-667.

[De Robertis, E. M.](#) (2006). Spemann's organizer and self-regulation in amphibian embryos. *Nat Rev Mol Cell Biol* 7, 296-302.

[Keller, R., Davidson, L. A., and Shook, D. R.](#) (2003). How we are shaped: the biomechanics of gastrulation. *Differentiation* 71, 171-205.

### Dominant negative approaches

\*[Amaya, E., Musci, T. J., and Kirschner, M. W.](#) (1991). Expression of a dominant negative mutant of the FGF receptor disrupts mesoderm formation in *Xenopus* embryos. *Cell* 66, 257-270.

[Hemmati-Brivanlou, A., and Melton, D. A.](#) (1992). A truncated activin receptor inhibits mesoderm induction and formation of axial structures in *Xenopus* embryos. *Nature* 359, 609-614.

### Antisense

[Heasman, J., Crawford, A., Goldstone, K., Garner-Hamrick, P., Gumbiner, B., McCrea, P., Kintner, C., Noro, C. Y., and Wylie, C.](#) (1994). Overexpression of cadherins and underexpression of beta-catenin inhibit dorsal mesoderm induction in early *Xenopus* embryos. *Cell* 79, 791-803.

[Xanthos, J. B., Kofron, M., Wylie, C., and Heasman, J.](#) (2001). Maternal VegT is the initiator of a molecular network specifying endoderm in *Xenopus laevis*. *Development* 128, 167-180.

### Timing and nature of signaling

\*[Wylie, C., Kofron, M., Payne, C., Anderson, R., Hosobuchi, M., Joseph, E., and Heasman, J.](#) (1996). Maternal beta-catenin establishes a 'dorsal signal' in early *Xenopus* embryos. *Development* 122, 2987-2996.

[Faure, S., Lee, M. A., Keller, T., ten Dijke, P., and Whitman, M.](#) (2000). Endogenous patterns of TGFbeta superfamily signaling during early *Xenopus* development. *Development* 127, 2917-2931.

[Lee, M. A., Heasman, J., and Whitman, M.](#) (2001). Timing of endogenous activin-like signals and regional specification of the *Xenopus* embryo. *Development* 128, 2939-2952.

[Agius, E., Oelgeschlager, M., Wessely, O., Kemp, C., and De Robertis, E. M.](#) (2000). Endodermal Nodal-related signals and mesoderm induction in *Xenopus*. *Development* 127, 1173-1183.

### Neural induction

[Lamb, T. M., Knecht, A. K., Smith, W. C., Stachel, S. E., Economides, A. N., Stahl, N., Yancopoulos, G. D., and Harland, R. M.](#) (1993). Neural induction by the secreted polypeptide noggin. *Science* 262, 713-718.

[Smith, W. C., and Harland, R. M.](#) (1992). Expression cloning of noggin, a new dorsalizing factor localized to the Spemann organizer in *Xenopus* embryos. *Cell* 70, 829-840.

\*[Kiecker, C., and Niehrs, C.](#) (2001). A morphogen gradient of Wnt/beta-catenin signalling regulates anteroposterior neural patterning in *Xenopus*. *Development* 128, 4189-4201.

\*[Khokha, M. K., Yeh, J., Grammer, T. C., and Harland, R. M.](#) (2005). Depletion of Three BMP Antagonists from Spemann's Organizer Leads to a Catastrophic Loss of Dorsal Structures. *Dev Cell* 8:401-11.

\*[Reversade, B., and De Robertis, E. M.](#) (2005). Regulation of ADMP and BMP2/4/7 at opposite embryonic poles generates a self-regulating morphogenetic field. *Cell* 123, 1147-60.

### Morphogenesis

[Davidson, L. A., and Keller, R. E.](#) (1999). Neural tube closure in *Xenopus laevis* involves medial migration, directed protrusive activity, cell intercalation and convergent extension. *Development* 126, 4547-4556.

[Haigo, S. L., Hildebrand, J. D., Harland, R. M., and Wallingford, J. B.](#) (2003). Shroom induces apical constriction and is required for hinge point formation during neural tube closure. *Curr Biol* 13, 2125-2137.

[Wallingford, J. B., Rowling, B. A., Vogeli, K. M., Rothbacher, U., Fraser, S. E., and Harland, R. M.](#) (2000). Dishevelled controls cell polarity during *Xenopus* gastrulation. *Nature* 405, 81-85.