

Stress-Regulated Targeting of the NKG2D Ligand Mult1 by a Membrane-Associated RING-CH Family E3 Ligase

Timothy J. Nice, Weiwen Deng, Laurent Coscoy, and David H. Raulet

NKG2D is a stimulatory receptor expressed by NK cells and some T cell subsets. Expression of the self-encoded ligands for NKG2D is presumably tightly regulated to prevent autoimmune disorders while allowing detection of infected cells and developing tumors. The NKG2D ligand Mult1 is regulated at multiple levels, with a final layer of regulation controlling protein stability. In this article, we report that Mult1 cell-surface expression was prevented by two closely related E3 ubiquitin ligases membrane-associated RING-CH (MARCH)4 and MARCH9, members of an E3 family that regulates other immunologically active proteins. Lysines within the cytoplasmic domain of Mult1 were essential for this repression by MARCH4 or MARCH9. Downregulation of Mult1 by MARCH9 was reversed by heat-shock treatment, which resulted in the dissociation of the two proteins and increased the amount of Mult1 at the cell surface. These results identify Mult1 as a target for the MARCH family of E3 ligases and show that induction of Mult1 in response to heat shock is due to regulated association with its E3 ligases. *The Journal of Immunology*, 2010, 185: 5369–5376.

NKG2D is a receptor expressed by NK cells and subsets of T cells. When engaged by cellular ligands, NKG2D initiates stimulatory signaling cascades by virtue of its association with the signaling adaptor molecules DAP10 and/or DAP12. Stimulation of immune cells through NKG2D was shown to be involved in protective responses to infections and tumors (1, 2), but it also can lead to autoimmunity (3). Therefore, the regulation of activating signals through NKG2D is extremely important to protect the host from pathogens and tumors while preventing the destruction of self-tissue. The NKG2D receptor is constitutively expressed on NK cells and some T cells, and the regulation of the NKG2D-mediated response occurs largely at the level of ligand expression.

The ligands for NKG2D are self-proteins that are absent or expressed at a low level on most normal tissues. There are nine known murine ligands (Rae1 α –Rae1 ϵ , H60a–H60c, and Mult1) and nine known human ligands (RAET1a–RAET1g [also known as UL-16 binding proteins], MICA, and MICB) that are all distantly related to MHC class I. However, because of their relatively rapid rate of evolution, it has not been possible to draw direct lines of homology between individual mouse and human ligands (4). Although induction of the ligands is known to occur on tumor cells and during certain infections, the mechanisms regulating this induction are still being characterized. Stress pathways, most notably the DNA damage and heat-shock pathways, were shown

to induce ligand expression on cell lines in vitro (5, 6). Although the DNA damage-response pathway was shown to regulate many of the NKG2D ligands similarly, the heat-shock pathway acts more selectively on a subset of ligands: MICA/B in humans and Mult1 in mice. However, heat-shock regulation differs in the latter two cases. Although MICA/B are thought to be regulated transcriptionally by the heat-shock response, Mult1 was shown to be induced as a consequence of heat-shock-mediated suppression of the high steady-state ubiquitination and turnover of Mult1 (7).

To better understand the molecular mechanism of posttranslational Mult1 regulation, it is important to identify E3 ubiquitin ligases that target Mult1. E3 ubiquitin ligases are, in most cases, noncatalytic scaffolds that bring a ubiquitin-bearing E2 ubiquitin-conjugating enzyme in close proximity to a target, allowing the transfer of ubiquitin to occur (8). Thus, target specificity of ubiquitination reactions are most often determined by an E3 ligase. The membrane-associated RING-CH (MARCH) family of E3 ligases (9, 10) represents a promising set of candidate Mult1 regulatory proteins for several reasons. MARCH proteins are transmembrane (TM) proteins, and most of their targets are also TM proteins, many with significance to the immune system (9, 11–15). Of particular interest is the finding that MHC class I expression is regulated by ubiquitination dependent on the MARCH4 and MARCH9 E3 ligases, as well as by virally encoded homologs of the MARCH proteins MIR1 and MIR2 (9, 16). MHC class II is also ubiquitinated by MARCH proteins MARCH1 and MARCH8, and regulation of this ubiquitination pathway is involved in the induction of MHC II expression on dendritic cells upon activation (11, 17).

Although the mechanism determining substrate recognition has not been determined for all of the endogenous MARCH proteins, the substrate specificity of MIR2 was shown, through domain swapping, to be dependent on their membrane-spanning regions (18). Because the TM domain of Mult1 is involved in its protein-level regulation (7), and Mult1 has homology to MHC class I, the MARCH proteins were prime candidates for Mult1 E3 ligases. This initial suspicion for a role of MARCH proteins in regulating Mult1 was bolstered by a recent report that the viral homolog MIR2 downregulates the human NKG2D ligands MICA and MICB (19). In this report, we provide direct evidence that MARCH4 and MARCH9, but not other MARCH proteins, can associate with Mult1 and suppress

Department of Molecular and Cell Biology and Cancer Research Laboratory, University of California at Berkeley, Berkeley, CA 94720

Received for publication January 26, 2010. Accepted for publication August 23, 2010.

Address correspondence and reprint requests to Dr. David H. Raulet, Department of Molecular and Cell Biology, University of California at Berkeley, 142 Life Sciences Addition #3200, Berkeley, CA 94720-3200. E-mail address: raulet@berkeley.edu

The online version of this article contains supplemental material.

Abbreviations used in this paper: C-M4, C1498-MARCH4 cells; C-M9, C1498-MARCH9 cells; C-MARCH9, C1498 cells transduced with mouse MARCH9; ctrl, untreated; HA, hemagglutinin; HS, heat shock; LAK, lymphokine activated killer; M9, MARCH9; MARCH, membrane-associated RING-CH; siRNA, small interfering RNA; TM, transmembrane; WT, wild-type.

Copyright © 2010 by The American Association of Immunologists, Inc. 0022-1767/10/\$16.00

Mult1 expression at the cell surface in a manner that depends on lysine residues in the Mult1 cytoplasmic domain and that can be reversed by heat shocking the cells.

Materials and Methods

Cells and heat shock

All cells were cultured in complete DMEM, consisting of DMEM (Life Technologies, Carlsbad, CA), 10% FCS (Omega Scientific, Tarzana, CA), 100 U/ml penicillin (Life Technologies), 100 µg/ml streptomycin (Life Technologies), 0.2 mg/ml glutamine (Sigma-Aldrich, St. Louis, MO), 10 µg/ml gentamicin sulfate (Lonza, Basel, Switzerland), 20 mM HEPES (Fisher BioReagents, Fair Lawn, NJ), and 50 µM 2-ME (EMD Chemicals, Gibbstown, NJ). Heat shock was performed by placing cells in a 45°C water bath for 30 min, followed by a 4-h incubation at 37°C. Dead cells were excluded from flow cytometry analysis through staining with 7-aminoactinomycin D.

Cytotoxicity assay

Lymphokine-activated killer (LAK) cells were prepared by culturing splenocytes for 4–5 d in 1000 U/ml IL-2. LAK cells were cocultured with ⁵¹Cr-labeled target cells in a standard 4-h [⁵¹Cr]-release assay (20).

Abs

Mult1 Ab (clone 237104; R&D Systems, Minneapolis, MN), H60a Ab (clone 205326; R&D Systems), hemagglutinin (HA) Ab (HA.11; Covance, Princeton, NJ), PE-conjugated goat F(ab')₂ fragment to rat IgG (Jackson ImmunoResearch Laboratories, West Grove, PA), PE-conjugated goat Ab to mouse IgG1 (Southern Biotechnology Associates, Birmingham, AL), PE-conjugated MHC class I (H-2D^b) Ab (clone 28-14-8; eBioscience, San Diego, CA), PE-conjugated H-2K^b Ab (clone AF6-88.5; eBioscience), and PE-conjugated ICAM-1 (clone YN1/1.7.4; eBioscience) were used for flow-cytometric analysis. Mult1 Ab (clone 1D6; kind gift of S. Jonjic, University of Rijeka, Rijeka, Croatia) (21) was used for immunoprecipitations and Western blots. Ubiquitin Ab (clone P4D1; Santa Cruz Biotechnology, Santa Cruz, CA), α-tubulin Ab (Calbiochem, San Diego, CA), myc Ab (clone 9E10; eBioscience), HA Ab (HA.11; Covance), HRP-conjugated goat anti-mouse IgG (Pierce, Rockford, IL), and peroxidase-conjugated goat anti-mouse L chain (Jackson ImmunoResearch Laboratories) were used for Western blots.

Cloning of MARCH cDNA

The human MARCH proteins were amplified from EST clones purchased from the American Type Culture Collection (Manassas, VA) and subcloned into pCGN-HA. Mouse MARCH4 and MARCH9 were amplified from mouse fibroblast cDNA using the following primers: BamHI-M4F, 5'-CCAAAG-GATCCCATGCTCATGCCCTGGGTG-3'; HindIII-M4R, 5'-CCAAAAA-GCTTTTCACACTGTGGTGACTCTCATGAC-3'; BamHI-M9F5', 5'-CCA-AAGGATCCGATGCTCAAGTCTCGGCTCCG-3'; M9R5', 5'-GAAGTA-GCAAAGCTCACAGCTCCATG-3'; M9F3', 5'-AAGTACCAGGTCCTG-CGATCAGCAC-3'; and HindIII-M9R3', 5'-CCAAAAAGCTTTCAGAG-TGTGGTGACCCCTCATG-3'. MARCH9 was cloned by assembling two amplified segments (5' end amplified from genomic DNA) due to difficulty in achieving full-length amplification through the GC-rich 5' end. Amplified mouse MARCH4 and MARCH9 products were subcloned into pCMV-TAG3a.

Quantitative RT-PCR

Total RNA was isolated using TRIzol LS (Invitrogen, Carlsbad, CA), followed by digestion of contaminating DNA using DNA-free (Ambion, Austin, TX), according to the manufacturer's protocol. RNA was reverse transcribed using Superscript III reverse transcriptase (Invitrogen), and the resulting cDNA was used for quantitative PCR. Duplicate or triplicate amplification mixtures were prepared with 0.01–1 µg cDNA, SYBR GreenER SuperMix (Invitrogen), and 200 nM forward and reverse primers and cycled using the ABI 7300 real-time PCR system. Cycling parameters used were 50°C for 2 min, 95°C for 10 min, and 40 cycles of 95°C for 15 s and 60°C for 60 s. The following primers were used: Mult1-5', 5'-CAATGCTCTGTCTCGGAA-3'; Mult1-3', 5'-CTGAACACGCTCA-GGCACT-3'; MARCH4-5', 5'-CTCTGGGAAGTAGCTTGGAC-3'; MARCH4-3', 5'-CTGAACCATCACAAACGGCATG-3'; MARCH9-5', 5'-TC-ATGGAGCTGTGAGCTTTG-3'; MARCH9-3', 5'-CAGGACGATAGCA-GCAATCT-3'; RPS29-5', 5'-AGCAGCTCTACTGGAGTACC-3'; and RPS29-3', 5'-AGGTCGCTTAGTCCAACCTTAATG-3'.

Retroviral constructs and transduction of cells

FLAG or HA-tagged Mult1 was cloned into the pMSCV2.2-IRES-GFP retroviral vector (22). Myc-tagged MARCH4 and MARCH9 were subcloned into pMSCV-IRES-Thy1.1. Mutants of Mult1 were made, as described (7). Retroviral supernatants were generated by cotransfecting 293T cells with plasmids encoding VSV gag/pol and env and pMSCV retroviral constructs using Lipofectamine 2000 (Invitrogen). Culture supernatants collected 48 h after transfection were added directly to actively proliferating cells, and transduced cells were sorted based on GFP or Thy1.1 expression.

Immunoprecipitation and Western blotting

Lysates were prepared by resuspending cell pellets in coimmunoprecipitation buffer (0.5% Nonidet P-40, 150 mM NaCl, 20 mM Tris [pH 7.4], 10% glycerol). Protease inhibitor mixture tablets (Roche, Basel, Switzerland) were added to the buffer immediately before use. When assessing ubiquitination of Mult1, 20 mM N-ethylmaleimide was included in the lysis buffer. Cells were lysed at 4°C for 30 min and cleared by centrifugation at 16,000 × g for 15 min. Lysates were precleared with protein G PLUS agarose beads (Santa Cruz Biotechnology). Precleared lysates were incubated with 1 µg Mult1 Ab (1D6) for 1 h, followed by incubation with protein G agarose beads overnight. Beads were washed three times in lysis buffer and boiled in SDS sample buffer for separation by SDS-PAGE. Samples resolved by SDS-PAGE were transferred to nitrocellulose membranes. Membranes were blocked in 5% milk and incubated with primary Abs, followed by HRP-coupled secondary Abs. Following incubation with Western lightning chemiluminescence reagent (PerkinElmer, Wellesley, MA), membranes were exposed to film and developed.

Transfection of small interfering RNA oligonucleotides

Relative levels of MARCH4 and MARCH9 were determined using the primers MARCH4+9-5' (5'-CAGTGGCAGCCAT-3') and MARCH4+9-3' (5'-ACTTTCCACTGCTGGTT-3'), followed by digestion with Sall, which selectively digests MARCH4 amplicons. Fifty thousand fibroblasts were transfected with 100 pmol small interfering RNA (siRNA) oligonucleotides using Lipofectamine 2000. The following oligonucleotides were used: GFP-22 (Qiagen, Valencia, CA), control oligonucleotide (Invitrogen), and three predesigned MARCH9 oligonucleotides (Invitrogen). Seventy-two hours later, cells were lifted and used for analysis of Mult1 surface expression and MARCH9 quantitative PCR.

Results

Screen of human MARCH proteins identifies MARCH4 and MARCH9 as Mult1 E3 ligases

The nine mammalian MARCH proteins that were originally described can be distinguished based on the number of TM domains they possess. MARCH1, MARCH2, MARCH3, MARCH4, MARCH8, and MARCH9 have two TMs and are most closely related to the viral homologs. Plasmids encoding HA-tagged versions of these two-TM MARCH proteins cloned from human cDNA were used to screen for the ability to regulate Mult1 expression. 293T cells were transiently cotransfected with individual human MARCH plasmids or control plasmid, together with a plasmid encoding Mult1 and IRES-GFP. Cells were stained 24 h later with Mult1 Abs, and transfected cells were selectively analyzed by gating on GFP-expressing cells. Strikingly, cotransfection of Mult1 with either of two highly related family members (human MARCH4 and human MARCH9) prevented Mult1 cell-surface expression (Fig. 1A). This effect was specific to human MARCH4 and human MARCH9, because no other MARCH family members that were tested (MARCH6, MARCH7, MARCH10 and MARCH11 were not examined) caused notable variations in Mult1 expression compared with control cells. Interestingly, these two proteins were the same MARCH proteins reported to downregulate MHC class I molecules (9).

To determine the role of the Mult1 cytoplasmic domain in MARCH9-mediated downregulation, 293T cells were cotransfected with Mult1 cytoplasmic domain mutants and human MARCH9 (Fig. 1B). Human MARCH9 strongly inhibited cell-




FIGURE 1. Screen of human MARCH proteins identifies MARCH4 and MARCH9 as Mult1 E3 ligases. 293T cells were cotransfected with a plasmid encoding WT Mult1-IRES-GFP and a plasmid encoding the indicated human MARCH proteins. *A* and *B*, GFP⁺ cells were analyzed 24 h after transfection by flow cytometry for Mult1 expression. Expression of GFP, which is encoded by the same bicistronic mRNA as Mult1, was similar in all of the samples, indicating comparable Mult1 gene expression. Shaded graphs represent staining of cells transfected with a plasmid encoding IRES-GFP only. *C*, Cells were lysed in coimmunoprecipitation buffer, followed by immunoprecipitation of Mult1 and detection of FLAG, HA, or ubiquitin by Western blotting. Similar results were obtained in two additional experiments.

surface expression of a Mult1 truncation mutant lacking aa 253–334 (the $\Delta 253$ mutant), showing that most of the cytoplasmic domain, containing four cytoplasmic lysine residues (K278, K280, K310, and K311), was not necessary for MARCH9-mediated downregulation. In contrast, human MARCH9 cotransfection did not impair cell-surface expression of a Mult1 truncation mutant (the $\Delta 239$ mutant) lacking a slightly larger segment, comprising amino acids 239–334, including two additional cytoplasmic lysine residues (K240 and K241) compared with the $\Delta 253$ mutant. The role of the cytoplasmic lysine residues was confirmed by the demonstration that the KR mutant, a full-length Mult1 in which six cytoplasmic lysine residues were substituted with arginine residues, was only slightly downregulated by human MARCH9. The marginal downregulation of the KR mutant by human MARCH9 was reproducible, however, suggesting that nonlysine residues in the cytoplasmic domain may be suboptimal targets for MARCH9-mediated ubiquitination, as was demonstrated in the case of ubiquitination targeted by the other MARCH family proteins, MIR1, MK3, and MARCH2 (23, 24). Together, these results show that the Mult1 cytoplasmic domain is critical for downregulation by MARCH9 and that lysines within this domain are required for this regulation.

The requirement for lysine residues within the cytoplasmic domain strongly suggested that the downregulation of Mult1 by MARCH9 involved ubiquitination. To confirm that Mult1 is ubiquitinated by MARCH4 and MARCH9, Mult1 was immunoprecipitated from transfected 293T cells, followed by detection of ubiquitin by Western blotting. High m.w. ubiquitin species were evident in immunoprecipitates from cells cotransfected with human MARCH4 or human MARCH9 and wild-type (WT) Mult1 but were notably less abundant when the KR or $\Delta 239$ Mult1 mutants were expressed (Fig. 1C). The fact that these ubiquitinated species were dependent on lysines in the Mult1 cytoplasmic domain, were absent in the vector-transfected cells, and were larger than Mult1 (55 kDa), strongly suggested that they represent ubiquitinated Mult1. Surprisingly, although WT but not KR Mult1 was ubiquitinated and depleted from the cell surface in the presence of

human MARCH4/9, there was no apparent decrease in total WT Mult1 protein levels seen by Western blot (Fig. 1C). This finding suggests that the downregulation of Mult1 in the presence of MARCH4/9 in 293T cells is due to sequestration in an intracellular compartment.

It remained possible that the effect of MARCH4/9 on Mult1 expression was indirect, so coimmunoprecipitation of the two proteins was performed to measure the extent of their physical interaction. Mult1 was immunoprecipitated from cotransfected cells, followed by Western blotting to detect HA-tagged human MARCH proteins (Fig. 1C). Human MARCH4 and human MARCH9 were coimmunoprecipitated with Mult1, showing that a physical interaction does occur. The interaction of Mult1 with the human MARCH proteins did not depend on the presence of lysines in the Mult1 cytoplasmic domain, because human MARCH4 and human MARCH9 were coimmunoprecipitated with the KR mutant to the same extent as with WT Mult1. In contrast, human MARCH4/9 failed to coimmunoprecipitate appreciably with the $\Delta 239$ mutant of Mult1, indicating that the cytoplasmic domain, independent of the cytoplasmic lysine residues, might be important for allowing interaction with MARCH4 or MARCH9. Together, these data show that MARCH4 and MARCH9 interact with Mult1 in a manner dependent on the Mult1 cytoplasmic domain, leading to its ubiquitination and reduced surface expression.

Mouse MARCH4/9 downregulate endogenous Mult1

Although the MARCH proteins are highly conserved between humans and mice, it was important to verify the results obtained with the human proteins using mouse MARCH4/9 in a more physiological experimental setting. Mouse MARCH4/9 were amplified from cDNA, N-terminally tagged with the myc epitope, and cloned into a MSCV retroviral expression vector. Retroviral transduction was used because it allowed stable ectopic expression at lower levels than transient transfection, thus minimizing potential nonspecific effects of forced expression. Two cell lines (C1498 and WEHI 7.1) were chosen for ectopic expression of mouse MARCH4/9 because they express endogenous Mult1.

Stable expression of mouse MARCH4 or MARCH9 in either of these cell lines led to decreased cell-surface expression of endogenous Mult1 (Fig. 2A, 2B), confirming that the mouse MARCH proteins downregulate Mult1. The downregulation of Mult1 was specific, because there was little or no downregulation of a distinct TM NKG2D ligand (H60a) that is normally expressed by WEHI 7.1 cells (Fig. 2C). In contrast, the endogenous Db class I MHC molecule was partially downregulated by mouse MARCH9 in C1498 cells, consistent with reports that transfected human MARCH4/9 downregulate MHC class I and ICAM-1 in human cells (Fig. 2D) (9, 12). In contrast, mouse MARCH9 failed to downregulate ICAM-1 or a distinct MHC I molecule, Kb, in transfected C1498 cells, suggesting selectivity in the effects of mouse MARCH proteins compared with human MARCH proteins (Fig. 2D). Although the possibility remains that mouse MARCH9 regulates other TM molecules not examined in this study in a more dramatic fashion, these results clearly show that targeting for downregulation by mouse MARCH9 is quite specific, and Mult1 is one of its targets.

MARCH9 expression inhibits lysis by NK cells

To determine whether MARCH proteins regulate the susceptibility of target cells to NK-mediated lysis, transduced C1498 cells were used as target cells for activated NK cells (LAK cells). C1498 cells transduced with mouse MARCH4 or mouse MARCH9 were killed significantly less well by LAK cells than by parental C1498 cells (Fig. 3A). A greater reduction in killing was observed with cells transduced with mouse MARCH9 compared with mouse MARCH4, which correlated with a greater reduction in Mult1 in the former cells (Fig. 3A). However, although mouse MARCH9-transduced cells were nearly devoid of cell-surface Mult1, the cells retained partial sensitivity to LAK cell killing, presumably as a result of the presence of additional stimulatory ligands for NK cells on C1498 cells that are not downregulated by MARCH9.

To determine the component of the reduced killing that was due to downregulation of Mult1, we compared killing by LAK cells prepared from WT or NKG2D knockout mice. Parental C1498 cells were killed significantly, although modestly, less well by NKG2D-deficient LAK cells than by WT LAK cells. Because C1498 cells express Mult1, but not other NKG2D ligands (data not shown), these data suggested that Mult1 contributed significantly to NK killing of C1498 cells but that ligands for other activating receptors also played a large role, as was true of most target cell lines studied




FIGURE 2. Mouse MARCH4 and MARCH9 downregulate endogenous Mult1. WEHI 7.1 (A, C) or C1498 (B, D) cells were transduced with retroviruses encoding mouse MARCH4 or mouse MARCH9. A coexpressed IRES-Thy1.1 cassette was used to mark transduced cells. Thy1.1⁺ cells were sorted and analyzed for expression of Mult1 (A, B), H60a (C), K^b, D^p, or ICAM-1 (D). Similar results were obtained in at least two additional experiments.




FIGURE 3. MARCH9 expression inhibits lysis by NK cells. A, Non-transduced C1498 cells or C1498 cells transduced with mouse MARCH4 or mouse MARCH9 were used as targets for activated NK cells (LAK cells) in a chromium-release assay. Target cells were analyzed for Mult1 expression by flow cytometry. B, C1498 or C1498-MARCH9 cells were used as targets for LAK cells prepared from WT or NKG2D knockout mice in a chromium release assay. Results are representative of at least two independent experiments. C-M4, C1498-MARCH4 cells; C-M9, C1498-MARCH9 cells. * $p < 0.05$.

elsewhere (20). In contrast, C1498-MARCH9 cells were killed equally well by WT or NKG2D-deficient LAK cells, showing that removal of Mult1 from the cell surface prevented NKG2D-dependent killing. However, it was notable that the reduction in killing accomplished by MARCH9 transduction was greater than the reduction in killing of parental C1498 cells resulting from NKG2D deficiency. Therefore, it is likely that MARCH9 transduction downregulated ligands for other activating NKR, in addition to downregulating Mult1. The identity of the other affected ligands in our analysis remains unclear, but there are numerous NK-activating ligands, some of which may remain to be identified. The affected ligands are probably not ICAM-1, CD48 (ligand for 2B4), or CD155 (ligand for DNAM-1), because we did not observe ICAM-I downregulation in MARCH9-transduced cells (Fig. 2D) and did not detect CD48 or CD155 expression on C1498 cells (data not shown). The possibility that NK stimulatory and inhibitory ligands other than Mult1 are also targets of MARCH9 was suggested by published data (9, 12, 15). Together, these data show that MARCH9-mediated downregulation of Mult1 and probably other activating ligands on C1498 cells inhibits functional recognition by NK cells.

MARCH9 and regulation of Mult1 by heat shock

In some cell types, including WEHI 7.1, a fibroblast cell line, and a subset of primary thymocytes, expression of Mult1 was shown to be inducible by heat shock (Fig. 4) (7). In contrast, WT C1498 cells normally express somewhat higher levels of cell-surface Mult1; no increase was observed after the cells were




FIGURE 4. MARCH9 and regulation of Mult1 by heat shock. WEHI 7.1 and C1498 cells, transduced or not with mMARCH9, were heat shocked at 45°C for 30 min, followed by a 4-h incubation at 37°C. Expression of Mult1 and D^b were analyzed by flow cytometry. Panels on the left and right are organized to show the relative effects of heat shock and MARCH9, respectively. Similar results were obtained in at least two additional experiments.

subjected to heat shock (Fig. 4). It is not known why Mult1 is not regulated by heat shock in C1498 cells, but transduction of mouse MARCH9 into C1498 cells greatly reduced the high basal level of Mult1, raising the possibility that MARCH9 restored a missing component in the pathway regulating Mult1. Indeed, when mouse MARCH9-transduced C1498 cells were heat shocked, substantial levels of Mult1 were restored to the cell surface, whereas little change in Mult1 expression was seen on parental C1498 cells (Fig. 4). These data suggest that the parental C1498 cells lack an E3 ligase for Mult1, and this defect is rescued by ectopic expression of MARCH9.

Cell-surface expression of the MHC class I molecule D^b was diminished by transduction of mouse MARCH9 in C1498 cells, so it seems to be a target of MARCH9. Strikingly, however, D^b, unlike Mult1, was not induced by heat shock in mouse MARCH9-transduced C1498 cells. In fact, for unknown reasons, heat shock caused a reduction in D^b levels at the cell surface in mouse MARCH9-transduced and parental C1498 cells (Fig. 4). Taken together, these data indicate that Mult1 downregulation by MARCH9 is reduced following heat shock, but MARCH9-mediated downregulation of other proteins, such as D^b, is not. The results suggest that heat shock-dependent regulation targets Mult1 rather than targeting the MARCH9 E3 ligase or other targets of MARCH9.

A dynamic association of Mult1 with MARCH9 underlies its regulation

The fact that D^b is not induced by heat shock in mouse MARCH9-transduced cells suggested that a global change in MARCH9 activity might not be the cause of Mult1 regulation. To test this possibility, RNA and protein levels were analyzed before and after heat shock. As expected, transcript levels for mouse MARCH9 were higher in mouse MARCH9-transduced cells compared with parental C1498 cells, but there was not a significant reduction in transcript levels following heat shock (Fig. 5A). Similarly, the transduced mouse MARCH9 protein, detected by Western blotting




FIGURE 5. A dynamic association of Mult1 with MARCH9 underlies its regulation. Cells treated as described in Fig. 4 were used for MARCH9 quantitative PCR (A), myc Western blot (B), or Mult1-MARCH9 coimmunoprecipitation (C). MARCH9 transcripts detected by quantitative PCR were normalized to RPS29. The band in the myc blot indicated by the asterisk (*) is a nonspecific background band, below which myc-MARCH9 is visible. Similar results were obtained in two additional experiments. C-MARCH9, C1498 cells transduced with mouse MARCH9; ctrl, untreated (no heat shock); HS, heat shock.

with Myc epitope Abs, was not noticeably depleted following heat shock (Fig. 5B). These data argue strongly against the possibility that heat shock stimulates Mult1 expression by reducing MARCH9 expression.

An alternative explanation for Mult1 regulation is that the interaction between Mult1 and MARCH9 is reduced or prevented following heat shock, thus preventing targeting of Mult1. To test this hypothesis, the extent of interaction between myc-tagged mouse MARCH9 and Mult1 was determined in lysates from C1498-MARCH9 cells, before or after heat shock. Similar amounts of Mult1 were detected in the immunoprecipitates, but the amount of mouse MARCH9 coimmunoprecipitated was substantially reduced (~5-fold) in heat-shocked cells (Fig. 5C, quantification not shown). This indicates that the association of mouse MARCH9 with Mult1 was reduced following heat shock. The reduced interaction of Mult1 and mouse MARCH9 correlated with the increased display of Mult1 on the cell surface (Fig. 4, top panels). These data are consistent with a dynamic Mult1-MARCH9 interaction and, because MARCH9 levels are not regulated by heat shock, suggest that a heat shock-sensitive event controls the extent of this interaction.

Analysis of endogenous MARCH9

Although transduction of mouse MARCH4 and mouse MARCH9 clearly resulted in suppression of Mult1, it was important to investigate the role of the endogenous proteins. No Abs against mouse MARCH4 or MARCH9 are available, so analysis of endogenous expression was limited to transcript levels. To determine the tissue distribution of these MARCH family members, cDNA from a panel of mouse tissues and cells was analyzed by quantitative RT-PCR for the presence of MARCH4 and MARCH9 transcripts (Fig. 6A). Transcripts for MARCH9 were represented




FIGURE 6. Analysis of endogenous MARCH9. *A*, RNA from the indicated cells and tissues was used for MARCH4 or MARCH9 quantitative RT-PCR, and relative quantity was determined by normalization to RPS29. An asterisk (*) indicates that transcript amounts were below the limit of detection. *B*, Nontransduced fibroblasts or fibroblasts expressing HA-tagged Mult1 were transfected with control oligonucleotides (solid line) or the indicated siRNA oligonucleotides (dashed line). Expression of endogenous Mult1, GFP, and HA-Mult1 was analyzed 72 h after transfection. *C*, RNA was extracted from the siRNA-transfected fibroblasts (*B*) for quantitative PCR analysis, and relative levels of Mult1 and MARCH9 transcripts were determined by normalizing to RPS29. Similar results were obtained in two additional experiments. M9, MARCH9.

in a broad range of tissues, but MARCH4 transcripts were more selectively expressed, most notably in the brain and lung. The tissue distribution observed was similar to published analysis of human MARCH4/9 expression (9), suggesting similar roles for these proteins in mice and humans.

To more directly address the relative expression level of MARCH9 and MARCH4, PCR primers were designed that amplified a homologous region of these highly related transcripts, including a distinguishing Sall restriction enzyme site that is unique to MARCH4. Sall digestion of the PCR products was used to estimate the relative abundance of transcripts encoding MARCH4 (cleaved) versus MARCH9 (uncleaved). An established mouse fibroblast line (5) was selected for this more detailed analysis because this line had been well characterized in terms of Mult1 regulation. The PCR product from fibroblast cDNA was only visible as uncleaved product, indicating that MARCH9 is the only transcript present in appreciable quantities in these cells (Supplemental Fig. 1). Therefore, MARCH9 could be specifically studied in these cells without the complication of appreciable coexpression of MARCH4.

Mult1 in fibroblasts is maintained at low levels under normal conditions as a result of ubiquitination and degradation. To determine whether the endogenous MARCH9 in these cells was responsible for low cell-surface expression, siRNA oligonucleotides targeting MARCH9 were used to deplete MARCH9 transcripts. Nontransduced fibroblasts or fibroblasts ectopically expressing WT Mult1 and GFP were transfected with GFP oligonucleotides, negative control oligonucleotides, or three different MARCH9 oligonucleotides. Although the GFP oligonucleotides efficiently inhibited GFP expression in $\geq 80\%$ of cells, demonstrating the high efficiency of transfection, none of the three MARCH9 oligonucleotides used for knockdown resulted in appreciable increases in endogenous or ectopic Mult1 expression (Fig. 6*B*). Analysis of RNA levels in these cells showed that the MARCH9 oligonucleotides were effective, reducing transcript amounts by 70–95% (Fig. 6*C*). Also, transfection of MARCH9 siRNA oligonucleotides into 3T3 cells ectopically expressing MARCH9 effectively restored Mult1 surface expression, further illustrating their effectiveness (Supplemental Fig. 2). To ensure that there were no off-target effects on Mult1 that could confound interpretation of the results, Mult1 transcripts were also quantified. Although one of the oligonucleotides (M9 #3) seemed to reduce Mult1 RNA levels in this and some other experiments, the MARCH9 oligonucleotide that resulted in the best specific knockdown (M9 #1) had no effect on Mult1 transcript amounts (Fig. 6*C*). It is unlikely that endogenous MARCH4 was responsible for downregulating Mult1 in the absence of MARCH9, because these cells expressed MARCH4 mRNA levels below the limit of detection (Fig. 6*A*). Moreover, no additional effect was observed when MARCH4 oligonucleotides were cotransfected with the MARCH9 oligonucleotides (T.J. Nice and D.H. Raulet, unpublished observations). The results argue that MARCH9 and MARCH4 are not necessary for the basal suppression of Mult1 expression in these cells, but they may carry out this function redundantly with another E3 ligase.

Discussion

The MARCH E3 ligases and their viral homologs are a recently described protein family. These proteins are distinguished from other RING domain-containing E3 ligases by their unique configuration of cysteines and a histidine. Interestingly, they are all TM proteins with at least two membrane-spanning segments, and most of their ubiquitination targets are TM proteins. Of the MARCH-targeted proteins that have been identified, most are TM mole-

cules within the immune system. Our results demonstrate that ectopic expression of two members of this family (MARCH4 and MARCH9) prevent the expression of Mult1, adding this NKG2D ligand to the list of MARCH-regulated proteins. Our data support the notion that endogenous MARCH proteins, particularly the two-TM MARCH proteins, play numerous roles in the regulation of immune functions by regulating the cell-surface display of proteins that activate immune responses.

MIR2, a homolog of the MARCH proteins encoded by KSHV, was recently shown to downregulate the human NKG2D ligands MICA and MICB (19). Because it is believed that MIR2 originated as a human MARCH gene that was captured by KSHV, it is plausible that human MARCH proteins may also regulate human NKG2D ligands. However, we failed to observe MICA or MICB downregulation in Jurkat cells transfected with any of the human two-TM MARCH cDNAs (T.J. Nice and D.H. Raulet, unpublished data). Hence, MIR2 and endogenous human MARCH proteins may have diverged in their capacity to target MICA/B. It remains possible that MICA or MICB is regulated by one of the MARCH proteins not tested in this study or that MARCH proteins regulate a member of the distinct ULBP family of NKG2D ligands. Alternatively, Jurkat cells may be subject to constitutive stress signals that prevent targeting of MICA/B by transfected MARCH proteins.

The interaction of MARCH4 or MARCH9 with Mult1 was specific in several respects. First, the other MARCH family members failed to appreciably downregulate Mult1 in the transfection experiments (Fig. 1A). Second, stable expression of mouse MARCH9 using retroviral transduction did not downregulate several other immunologically relevant cell-surface proteins, including several potential MARCH targets, such as ICAM-1, H60a, and K^b. In contrast, the class I molecule D^b, another potential target, was downregulated, although not to the same extent as Mult1. These demonstrations of specificity argue against a non-specific effect of overexpression. In fact, the level of MARCH9 RNA in transduced C1498 cells (Fig. 5A) was only ~5-fold greater than levels seen in tissues with the most abundant MARCH9 RNA (e.g., brain). Given the likely heterogeneity of expression in a tissue, the level of MARCH9 mRNA in individual cells in the tissue may be substantially higher. These considerations suggest that MARCH9 is not grossly overexpressed in C1498-MARCH9 cells.

Although ectopically expressed MARCH4/9 led to downregulation of Mult1 expression, MARCH4 and MARCH9 are probably not the exclusive E3 ligases for Mult1, because knockdown of MARCH9 in fibroblasts that did not express detectable levels of MARCH4 transcript had no effect on Mult1 expression. The most likely explanation for this observation is redundancy of MARCH4/9 with unidentified cellular E3 ligases. These could be untested MARCH family members or non-MARCH E3 ligases. Notably, MARCH4/9 transcripts were more abundant in several tissues (e.g., brain, lung, and kidney; Fig. 6A) than in the cell lines analyzed, so MARCH4/9 may have a more prominent role in Mult1 regulation in the context of those tissues.

Although expression of MARCH4 or MARCH9 in 293T cells and C1498 cells led to a strong downregulation of Mult1 from the cell surface, Western blotting showed that they caused no (Figs. 1C, 4C), or at most a 2-fold, reduction (T.J. Nice, W. Deng, and D.H. Raulet, unpublished data) in total Mult1 amounts in these cells. These observations suggest that Mult1 downregulation by MARCH4/9 was largely due to sequestration within the cell. MHC class II regulation by MARCH1/8 was also shown to be partially due to sequestration in intracellular compartments, although some degradation was also observed (17). In contrast, downregulation of Mult1 mediated by endogenous E3 ligases in fibroblasts resulted

in very substantial reductions in total cellular Mult1 (7). Hence, in distinct circumstances, E3 ligases may prevent Mult1 cell-surface expression by degrading the protein or sequestering it in an intracellular compartment.

Interestingly, Mult1 was induced by heat shock in fibroblasts, where downregulation was not dependent on MARCH9, and in C1498-MARCH9 cells, where MARCH9 is the exclusive relevant E3 ligase. In terms of the mechanism of heat-shock regulation, these findings suggest that Mult1, rather than the E3 ligases, is targeted. This idea is supported by the observations that MARCH9 levels in C1498-MARCH9 cells did not change in response to heat shock (Fig. 5), and heat shock did not induce expression of the distinct MARCH9-targeted protein D^b (Fig. 4). A motif within the Mult1 cytoplasmic domain may be targeted by this stress response, in a manner that controls the interaction of Mult1 with its E3 ligases. Indeed, the Mult1 cytoplasmic tail is required for physical association with MARCH4/9; therefore, it may also be important for the regulation of their interaction following heat shock. This interaction could be disrupted by the association of a regulatory molecule with the cytoplasmic tail that prevents E3 binding and/or ubiquitination or that targets Mult1 to a compartment inaccessible to the E3 ligase. Alternatively, heat shock may repress a regulator that facilitates E3 binding and/or ubiquitination of Mult1 but not other targets of MARCH4/9. Further analysis is required to identify critical motifs within the cytoplasmic domain and how they regulate E3 ligase binding in response to heat shock.

Disclosures

The authors have no financial conflicts of interest.

References

- Guerra, N., Y. X. Tan, N. T. Joncker, A. Choy, F. Gallardo, N. Xiong, S. Knoblaugh, D. Cado, N. M. Greenberg, N. R. Greenberg, and D. H. Raulet. 2008. NKG2D-deficient mice are defective in tumor surveillance in models of spontaneous malignancy. [Published erratum appears in 2008 *Immunity* 28: 723.] *Immunity* 28: 571–580.
- Jonjić, S., M. Babić, B. Polić, and A. Krmpotić. 2008. Immune evasion of natural killer cells by viruses. *Curr. Opin. Immunol.* 20: 30–38.
- Ogasawara, K., J. A. Hamerman, H. Hsin, S. Chikuma, H. Bour-Jordan, T. Chen, T. Pertel, C. Carnaud, J. A. Bluestone, and L. L. Lanier. 2003. Impairment of NK cell function by NKG2D modulation in NOD mice. *Immunity* 18: 41–51.
- Raulet, D. H. 2003. Roles of the NKG2D immunoreceptor and its ligands. *Nat. Rev. Immunol.* 3: 781–790.
- Gasser, S., S. Orsulic, E. J. Brown, and D. H. Raulet. 2005. The DNA damage pathway regulates innate immune system ligands of the NKG2D receptor. *Nature* 436: 1186–1190.
- Groh, V., S. Bahram, S. Bauer, A. Herman, M. Beauchamp, and T. Spies. 1996. Cell stress-regulated human major histocompatibility complex class I gene expressed in gastrointestinal epithelium. *Proc. Natl. Acad. Sci. USA* 93: 12445–12450.
- Nice, T. J., L. Coscoy, and D. H. Raulet. 2009. Posttranslational regulation of the NKG2D ligand Mult1 in response to cell stress. *J. Exp. Med.* 206: 287–298.
- Pickart, C. M. 2001. Mechanisms underlying ubiquitination. *Annu. Rev. Biochem.* 70: 503–533.
- Bartee, E., M. Mansouri, B. T. Hovey Nerenberg, K. Gouveia, and K. Früh. 2004. Downregulation of major histocompatibility complex class I by human ubiquitin ligases related to viral immune evasion proteins. *J. Virol.* 78: 1109–1120.
- Ohmura-Hoshino, M., E. Goto, Y. Matsuki, M. Aoki, M. Mito, M. Uematsu, H. Hotta, and S. Ishido. 2006. A novel family of membrane-bound E3 ubiquitin ligases. *J. Biochem.* 140: 147–154.
- De Gassart, A., V. Camosseto, J. Thibodeau, M. Ceppi, N. Catalan, P. Pierre, and E. Gatti. 2008. MHC class II stabilization at the surface of human dendritic cells is the result of maturation-dependent MARCH I down-regulation. *Proc. Natl. Acad. Sci. USA* 105: 3491–3496.
- Hoer, S., L. Smith, and P. J. Lehner. 2007. MARCH-IX mediates ubiquitination and downregulation of ICAM-1. *FEBS Lett.* 581: 45–51.
- Lehner, P. J., S. Hoer, R. Dodd, and L. M. Duncan. 2005. Downregulation of cell surface receptors by the K3 family of viral and cellular ubiquitin E3 ligases. *Immunol. Rev.* 207: 112–125.
- Nakamura, N., H. Fukuda, A. Kato, and S. Hirose. 2005. MARCH-II is a syntaxin-6-binding protein involved in endosomal trafficking. *Mol. Biol. Cell* 16: 1696–1710.

15. Hör, S., T. Ziv, A. Admon, and P. J. Lehner. 2009. Stable isotope labeling by amino acids in cell culture and differential plasma membrane proteome quantitation identify new substrates for the MARCH9 transmembrane E3 ligase. *Mol. Cell. Proteomics* 8: 1959–1971.
16. Coscoy, L., D. J. Sanchez, and D. Ganem. 2001. A novel class of herpesvirus-encoded membrane-bound E3 ubiquitin ligases regulates endocytosis of proteins involved in immune recognition. *J. Cell Biol.* 155: 1265–1273.
17. Shin, J. S., M. Ebersold, M. Pypaert, L. Delamarre, A. Hartley, and I. Mellman. 2006. Surface expression of MHC class II in dendritic cells is controlled by regulated ubiquitination. *Nature* 444: 115–118.
18. Sanchez, D. J., L. Coscoy, and D. Ganem. 2002. Functional organization of MIR2, a novel viral regulator of selective endocytosis. *J. Biol. Chem.* 277: 6124–6130.
19. Thomas, M., J. M. Boname, S. Field, S. Nejentsev, M. Salio, V. Cerundolo, M. Wills, and P. J. Lehner. 2008. Down-regulation of NKG2D and Nkp80 ligands by Kaposi's sarcoma-associated herpesvirus K5 protects against NK cell cytotoxicity. *Proc. Natl. Acad. Sci. USA* 105: 1656–1661.
20. Jamieson, A. M., A. Diefenbach, C. W. McMahon, N. Xiong, J. R. Carlyle, and D. H. Raulet. 2002. The role of the NKG2D immunoreceptor in immune cell activation and natural killing. *Immunity* 17: 19–29.
21. Krmpotic, A., M. Hasan, A. Loewendorf, T. Saulig, A. Halenius, T. Lenac, B. Polic, I. Bubic, A. Kriegeskorte, E. Pernjak-Pugel, et al. 2005. NK cell activation through the NKG2D ligand MULT-1 is selectively prevented by the glycoprotein encoded by mouse cytomegalovirus gene m145. *J. Exp. Med.* 201: 211–220.
22. Ranganath, S., W. Ouyang, D. Bhattacharya, W. C. Sha, A. Grupe, G. Peltz, and K. M. Murphy. 1998. GATA-3-dependent enhancer activity in IL-4 gene regulation. *J. Immunol.* 161: 3822–3826.
23. Cadwell, K., and L. Coscoy. 2005. Ubiquitination on nonlysine residues by a viral E3 ubiquitin ligase. *Science* 309: 127–130.
24. Wang, X., L. Lybarger, R. Connors, M. R. Harris, and T. H. Hansen. 2004. Model for the interaction of gammaherpesvirus 68 RING-CH finger protein mK3 with major histocompatibility complex class I and the peptide-loading complex. *J. Virol.* 78: 8673–8686.